



AP PHYSICS I
Chapter 9 Practice

Name: _____ Period: _____ Date: _____

1. A 2.0 kg frictionless cart is moving at a constant speed of 0.22 m/s to the right on a horizontal surface, as shown below, when it collides with a second cart of undetermined mass m that is initially at rest. As a result of the collision, the second cart acquires a speed of 1.6 m/s to the right. The graph of the velocity of the cart as a function of time t is shown below. Assume that friction is negligible before, during, and after the collision.



- a) Determine the cart's momentum before the collision.
- b) Determine the magnitude of the impulse during the collision.
- c) Determine the mass of the second cart.



2. Several students are riding in bumper cars at an amusement park. The combined mass of car A and its occupants is 250 kg. The combined mass of car B and its occupants is 200 kg. Car A is 15 m away from car B and moving to the right at 2.0 m/s, as shown, when the driver decides to bump into car B, which is at rest.

(a) Car A accelerates at 1.5 m/s^2 to a speed of 5.0 m/s and then continues at constant velocity until it strikes car B. Calculate the total time for car A to travel the 15 m.

(b) After the collision, car B moves to the right at a speed of 4.8 m/s.

i. Calculate the speed of car A after the collision.



ii. Indicate the direction of motion of car A after the collision.

_____ To the left _____ To the right _____ None; car A is at rest.

(c) Is this an elastic collision?

_____ Yes _____ No

Justify your answer.



3. A 2-kilogram block initially hangs at rest at the end of two 1-meter strings of negligible mass as shown on the left diagram above. A 0.003-kilogram bullet, moving horizontally with a speed of 1000 meters per second, strikes the block and becomes embedded in it. After the collision, the bullet/block combination swings upward, but does not rotate.

- (a) Calculate the speed of the bullet/block combination just after the collision.
- (b) Calculate the ratio of the initial kinetic energy of the bullet to the kinetic energy of the bullet/block combination immediately after the collision.
- (c) Calculate the maximum vertical height above the initial rest position reached by the bullet/block combination.